

Adjustable Gastric Banding (LAP-BAND™ or REALIZE™ Band) Patient Discharge Information

Follow-up appointments:

- Call Dr. Sandor's office (781-279-1123) once you get home to arrange a follow-up appointment for about 7 days after your surgery..
- You should already have a nutrition appointment scheduled for 3 weeks from the date of your surgery. If you do not have one scheduled, call (781) 756-2615 (for Winchester Hospital) or (781) 391-2121 (for Hallmark Health) as soon as possible to arrange.
- An appointment with your primary care physician about 4-6 weeks after surgery is advised, particularly, if you are on diabetic medication or blood pressure medication.

When to call the doctors office:

- Call if you have any fever/temperature > 101 degrees; any drainage from your incisions which is purulent (pus like/tan/thick); or warmth and redness surrounding the incisions.
- Call if you are having continuous nausea and vomiting and are unable to keep fluids down.
- Call if you have increasing abdominal pain or pain not responding to pain medication.
- Call for any questions or concerns that you may have.

What you can do:

- Move around as much as you want. Walking is good. Stairs are okay.
- You may shower. Get plenty of rest.
- Continue with Stage 3 Bariatric Diet and other fluids as per the handout (until you see nutrition – about 2-3 weeks). **The recommended daily protein volume is 70 g, which equals to 7 scoops of protein powder if you use the Metagenics Bariatric Meal product.**
- Use pain medication as needed. When able, may switch to Tylenol for decreasing pain.

What you cannot do:

No driving while taking pain medication. Once you are off of pain medication, your reaction times are good and you are able to move and change your position readily, you may resume driving. No jumping, running or heavy lifting: nothing > 20 pounds for 4 weeks after surgery.

What to expect:

You may have some days when you are not able to take in all of the liquid requirements of the diet. **Do not force the liquids** or try to play catch up.

Constipation may be a problem. If you have not had a bowel movement a couple of days after discharge from the hospital, you may use milk of magnesia. If no response you may try a Dulcolax suppository or a fleet enema. If constipation is an ongoing issue, you may require fiber supplementation (Metamucil) or a stool softener such as Colace, which is available over the counter.